

PROPAGATING FRONTS FOR VISCOUS SCALAR CONSERVATION LAW COUPLED WITH AN ELLIPTIC EQUATION

GIADA CIANFARANI CARNEVALE, CORRADO LATTANZIO,
AND CORRADO MASCIA

ABSTRACT. Motivated by radiation hydrodynamics, we analyse a 2×2 system consisting of a one-dimensional viscous conservation law with strictly convex flux –the viscous Burgers’ equation being a paradigmatic example– coupled with an elliptic equation. In the regime of small viscosity and for large shocks, namely when the profile of the corresponding underlying inviscid model undergoes a discontinuity –usually called *sub-shock*– it is proved the existence of a smooth propagating front, which regularises the jump of the corresponding inviscid equation. The proof is based on *Geometric Singular Perturbation Theory* (GSPT) as introduced in the pioneering work of Fenichel [5] and subsequently developed by Szmolyan [19]. In addition, the case of small shocks and large viscosity is also addressed via a standard bifurcation argument.

KEYWORDS. parabolic-elliptic system, traveling waves, singular perturbation theory.

2020 MSC CLASSIFICATION. 35C07, 34E15, 35B25, 76N30, 35A24.

1. INTRODUCTION

The dynamics of a gas in presence of radiation can be described by the classical compressible Euler equations with an additional term in the energy balance modelling the radiation effects (see [16, 21]). The extra state variable, describing the intensity of radiation, is often modelled positing that it obeys to an elliptic equation, leading to the hyperbolic–elliptic system

$$\left\{ \begin{array}{l} \partial_t \rho + \partial_x(\rho u) = 0 \\ \partial_t(\rho u) + \partial_x(\rho u^2 + p) = 0 \\ \partial_t(\rho E) + \partial_x(\rho E u + p u - \kappa \partial_x n) = 0 \\ -\partial_{xx} n + \tau \{n - g(\theta)\} = 0 \end{array} \right. \quad (1.1) \quad \boxed{\text{generalrad}}$$

where ρ denotes density, u velocity, p pressure, E specific total energy and θ temperature to be linked by some appropriate constitutive relations. The classical Euler system is then coupled through the term $\kappa \partial_x n$ to an additional quantity driven by the variable n , describing the average of the photon density by means of the positive parameters κ and τ , and the function g , who is usually assumed to have a power-like form. Such model is obtained in the non relativistic limit of a corresponding hyperbolic–kinetic system

where the variable describing the photon density obeys to a transport equation with interaction kernel given by the Stefan–Boltzmann law (see [1, 6, 14] for further details).

A simplified model for radiation dynamics is the 2×2 system for the scalars unknowns u and v having the specific form

$$\begin{cases} \partial_t u + \partial_x \left(\frac{1}{2} u^2 \right) = \partial_x v \\ v - \partial_{xx} v = \partial_x u, \end{cases} \quad (1.2) \quad \boxed{\text{Hamoriginal}}$$

which is known as *Hamer model* for radiating gas, see [7] (setting $q := -v$). The inviscid scalar conservation law is augmented with a scalar elliptic equation, mimicking the coupling present in system (1.1). The well-posedness of (1.2) is proved in [10] by adding a vanishing viscosity term, thus ending up in a parabolic-elliptic system (see also [2, 20]).

In the analysis of both systems (1.1) and (1.2), an intriguing issue is the study of existence and stability of shock profiles. Specifically, the weak dissipation properties –due to the coupling with the elliptic equation– give rise to the existence of sub-jumps for sufficiently strong shocks. Correspondingly, an increasing amount of regularity of the profile emerges as the magnitude of the shock decreases.

Such problem has been addressed in various cases, that is for the “scalar” models –a single conservation law with general flux functions coupled with the elliptic equation– both for small and regular profiles [18], and for possibly large and discontinuous ones [8, 11, 12]. The cases of systems has been addressed as well, and in particular for the specific Euler model with radiation effects (1.1) in [13] for weak (regular) profiles, and in [3, 15] for strong (discontinuous) shocks. The case of general hyperbolic–elliptic systems for small and large shocks is investigated in [11] for linear coupling and in [12] for nonlinear coupling.

Here, we are interested in exhibiting how the presence of a viscosity term in the simplified Hamer model (1.2) modifies the existence and regularity properties of the shock profiles. Therefore, in the sequel we consider the following regularised version of system (1.2)

$$\begin{cases} \partial_t u + \partial_x f(u) - \epsilon \partial_{xx} u = \partial_x v \\ v - \partial_{xx} v = \partial_x g(u), \end{cases} \quad (1.3) \quad \boxed{\text{genintro}}$$

where the flux function $f \in C^2(\mathbb{R})$ is (uniformly) convex and the coupling function $g \in C^1(\mathbb{R})$ is (strictly) increasing, that is

$$d^2 f(u) > 0 \quad \text{and} \quad dg(u) > 0 \quad \text{for any } u \in \mathbb{R}, \quad (1.4) \quad \boxed{\text{hypFG}}$$

where d denotes the derivative with respect to u . The corresponding *one-field equation* for the unknown u is

$$\partial_t u + \partial_x f(u) - \partial_{xx}(\epsilon u + g(u)) = \partial_{xx} \{ \partial_t u + \partial_x f(u) - \epsilon \partial_{xx} u \} \quad (1.5) \quad \boxed{\text{onefield}}$$

The main topic of this paper is to investigate the existence of shock profiles for model (1.3) –equivalently, for (1.5)– in the specific case of end states leading to the presence of a sub-shock in the inviscid model (see [8, 11, 12]).

For completeness, let us start with a classical definition.

Definition 1.1. *Given the states $u_{\pm} \in \mathbb{R}$, a **propagating front** for the hyperbolic-elliptic system (1.3) is a triple $(u, v; c)$ where (u, v) is a travelling wave solution $(u, v) = (u, v)(x - ct)$ satisfying the asymptotic conditions $(u, v)(\pm\infty) = (u_{\pm}, 0)$ and c is a given constant.*

*The couple (u, v) is called the **profile** and the parameter c is the **speed**.*

Incidentally, we are denoting with the same symbol both the general solution to (1.3), and the specific propagating front. Being this manuscript devoted exclusively to the existence of fronts, we are confident that this will not generate confusion.

The asymptotic states (u_{\pm}, v_{\pm}) are forced to be equilibria of (1.3), i.e. $u_{\pm} \in \mathbb{R}$ with $u_+ \neq u_-$ and $v_{\pm} = 0$. Consequently, the speed c is forced to satisfy the *Rankine–Hugoniot relation*

$$c = \frac{f(u_+) - f(u_-)}{u_+ - u_-}, \quad (1.6) \quad \boxed{\text{RH}}$$

as it is readily seen by integrating in \mathbb{R} the ordinary differential equation for the profile (u, v) . A standard assumption in the context of scalar conservation laws with convex flux is the *Lax condition*

$$df(u_+) < c < df(u_-), \quad (1.7) \quad \boxed{\text{Lax}}$$

guaranteeing that the jump from u_- to u_+ satisfies the entropy condition. Given the Rankine–Hugoniot condition (1.6) and the convexity assumption on the flux f , inequalities (1.7) are satisfied if and only if $u_+ < u_-$.

Next, we state the two main theorems of our work. The first one is based on a *singular perturbation* approach relative to the (small) parameter ϵ .

thm:main1

Theorem 1.2. *Assume (1.4)-(1.6)-(1.7). Moreover, let u_+, u_- be such that the inviscid profile undergoes a sub-shock in the u variable. Then, for $\epsilon > 0$ sufficiently small, the parabolic-elliptic system (1.3) support propagating fronts with speed c given by the Rankine-Hugoniot relation (1.6).*

This result is obtained in the case of small viscosity coefficient ϵ and for sufficiently large shocks, namely when the corresponding **inviscid** profile experiences a discontinuity, by taking advantage of the Fenichel’s *Geometric*

Singular Perturbation Theory, in what follows shortened as GSPT (for details, see [5, 19] and the Appendix at the end of the paper). The condition which locates the sub-shock in the profiles for the inviscid models agrees with the one needed to apply the aforementioned theory, leading to the existence of the connection which smoothes out the discontinuity.

Our second main result, proved in Section 3, is based on a direct application of bifurcation theory, where we consider fixed $O(1)$ viscosity and sufficiently small shocks.

thm:main3

Theorem 1.3. *Assume (1.4)-(1.6)-(1.7). Moreover, let u_+, u_- be such that $|u_+ - u_-|$ is sufficiently small. Then for any fixed $\epsilon > 0$ the parabolic-elliptic system (1.3) supports propagating fronts with speed c given by the Rankine-Hugoniot relation (1.6).*

The paper includes a final Appendix A with, the main general results of GSPT needed in the present analysis.

Notations. For readers' convenience, we list (some) symbols used in the text:

- x and $y = x/\epsilon$ denotes the slow and fast variable, respectively;
- $\mathbf{u} = (u, v, w)$ slow/fast variables for $\epsilon \geq 0$;
- f' and \dot{f} denotes the derivatives of the function f with respect to the slow and the fast variable, respectively;
- $\mathcal{S} := \{\mathbf{u} : F(\mathbf{u}) = 0\}$ slow manifold with its subsets $\mathcal{S}_*, \mathcal{S}_\pm$;
- $\mathbf{v} = (u, v, w, \epsilon)$ slow/fast variables;
- $\pi^{\mathcal{S}}$ projection onto the surface \mathcal{S}
- $\mathcal{C}, \mathcal{C}^s, \mathcal{C}^u$ center/center-stable/center-unstable manifolds for equilibrium point of limiting fast system
- $\mathcal{F}^{s,u} = \{\Psi^{s,u}(\mathbf{v}) : \mathbf{v} \in \mathcal{C}^{s,u}\}$ family of stable/unstable manifolds for $\mathcal{C}^{s,u}$
- $W^{s,u}$ stable/unstable manifolds of critical points for reduced slow system

However, all the notations will be restated at their first occurrence.

sec:genconvnon

2. GENERAL STRICTLY CONVEX CASE AND NONLINEAR COUPLING

To start with, we rephrase the existence of a propagating front in the formalism of singular perturbation theory.

Since f is convex, the function df is strictly increasing and, thanks to (1.7), there exists a unique point u_* such that $df(u_*) = c$. Hence, setting

$$f_c(u) := f(u) - cu. \quad (2.1) \quad \text{deficit}$$

there holds $df_c(u_*) = 0$. The function f_c describe the flux in a frame that is co-moving with the propagating front. The value $f_c(u_+)$ coincides with $f_c(u_-)$ as a consequence of the Rankine-Hugoniot condition (1.6) and we use the notation $f_c(u_\pm)$ to stress this coincidence. In addition, the Lax condition becomes

$$df_c(u_+) < 0 < df_c(u_-). \quad (2.2) \quad \text{LaxBis}$$

For traveling wave solutions, system (1.3) rewrites as

$$\epsilon u'' = \{f_c(u) - v\}', \quad v'' = v - g(u)',$$

where we have used the notation $' = d/dx$. Integrating the first equation from $-\infty$ to x , and setting $w := g(u) - v'$, we obtain the *slow system*

$$\begin{cases} \epsilon u' = F(u, v, w) := f_c(u) - f_c(u_{\pm}) - v, \\ v' = G(u, v, w) := w - g(u), \\ w' = H(u, v, w) := v, \end{cases} \quad (2.3) \quad \boxed{\text{SLOW SYSTEM}}$$

which is a special case belonging to the class of *singular perturbed problems*. The adjective *singular* refers to the presence of the parameter ϵ multiplying first order derivatives of some state variables and assumed to be small (and positive). A significant regime is then obtained as $\epsilon \rightarrow 0$ giving raise to the reduced limit

$$\begin{cases} v = f_c(u) - f_c(u_{\pm}), \\ v' = w - g(u), \\ w' = v. \end{cases} \quad (2.4) \quad \boxed{\text{SLOW SYSTEM } 0}$$

which is called *limiting slow system*. The first equality in (2.4) can be regarded as an additional constraint along the dynamics, linking the value of the variable u with the *slow variables* (v, w) . As in the present case, the relation $F(u, v, w) = 0$ is not invertible with respect to u and the use of appropriate restrictions have to be considered.

After the rescaling $y := x/\epsilon$, we get the so-called *fast system*

$$\begin{cases} \dot{u} = F(u, v, w), \\ \dot{v} = \epsilon G(u, v, w), \\ \dot{w} = \epsilon H(u, v, w). \end{cases}$$

with the notation $\dot{} = d/dy = \epsilon d/dx$. In compact form, setting $\mathbf{u} = (u, v, w)$, the above system can be rewritten as

$$\dot{\mathbf{u}} = \mathbf{F}_{\epsilon}(\mathbf{u}) := (F(\mathbf{u}), \epsilon G(\mathbf{u}), \epsilon H(\mathbf{u})). \quad (2.5) \quad \boxed{\text{FAST SYSTEM}}$$

As $\epsilon \rightarrow 0^+$ in (2.5), we deduce the *limiting fast system*

$$\begin{cases} \dot{u} = f_c(u) - f_c(u_{\pm}) - v, \\ \dot{v} = 0, \\ \dot{w} = 0. \end{cases} \quad (2.6) \quad \boxed{\text{FAST SYSTEM } 0}$$

The variable u is said to be the *fast variable* of the system.

From now on, we are confronted with the problem of rigorously establishing the existence of a heteroclinic orbit connecting at $\mp\infty$ the critical points of (2.3) –or, equivalently, (2.5)– which are

$$\mathbf{u}_{\pm} = (u_{\pm}, v_{\pm}, w_{\pm}) := (u_{\pm}, 0, g(u_{\pm})).$$

under the assumptions (1.4)–(2.2). To this aim, we follow the classical singular perturbation approach, consisting in the separate study of dynamical systems (2.3)–(2.5) in the limiting slow and fast regimes described by (2.4)–(2.6), respectively.

Let $\mathcal{S} := \{\mathbf{u} \in \mathbb{R}^3 : F(\mathbf{u}) = 0\}$ be the *slow manifold* of critical points for (2.6) and let \mathcal{S}_* be the open subset of the slow manifold \mathcal{S} defined by

$$\begin{aligned}\mathcal{S}_* &:= \{\mathbf{u} \in \mathcal{S} : \partial_u F(\mathbf{u}) \neq 0\} \\ &= \mathcal{S} \setminus \{(u_*, f_c(u_*) - f_c(u_\pm), 0) + t\mathbf{e}_3 : t \in \mathbb{R}\},\end{aligned}$$

Since the restrictions of the function $\partial_u F$ in the open halflines $(-\infty, u_*)$ and $(u_*, +\infty)$ are invertible, system (2.4) reduces to a two dimensional dynamical system for the slow variables (v, w) . Denoting such inverse functions with the symbols h_\pm , we can split the subset \mathcal{S}_* as the (disjoint) union

$$\mathcal{S}_* = \mathcal{S}_- \cup \mathcal{S}_+$$

where

$$\begin{aligned}\mathcal{S}_- &:= \{\mathbf{u} \in \mathcal{S}_* : \partial_u F(\mathbf{u}) > 0\} \\ &= \{(u, v, w) \in \mathbb{R}^3 : v \in (f_c(u_*) - f_c(u_\pm), +\infty), u = h_-(v)\}, \\ \mathcal{S}_+ &:= \{\mathbf{u} \in \mathcal{S}_* : \partial_u F(\mathbf{u}) < 0\} \\ &= \{(u, v, w) \in \mathbb{R}^3 : v \in (f_c(u_*) - f_c(u_\pm), +\infty), u = h_+(v)\}.\end{aligned}$$

By Lax condition (2.2), there holds $\mathbf{u}_- \in \mathcal{S}_-$ and $\mathbf{u}_+ \in \mathcal{S}_+$.

The analysis for the slow system (2.3) exhibits the existence of a discontinuous propagating front for $|u_+ - u_-|$ sufficiently large (in the case of the Hamer model (1.2), the critical threshold is $\sqrt{2}$, as shown in [8]). Specifically, we are interested in showing how the presence of viscosity in the first equation of (1.3) restores the regularity of the profile by smoothing out the internal jump of the shock solution. This task can be accomplished by working on the fast system at $\epsilon = 0$ from [8, 11, 12] and, then, taking advantage from the results in [19] to perform a rigorous matching.

2.1. Splitting consistency. The first step consists in the analysis of the linearization at \mathbf{u}_\pm of the fast system (2.5) for $\epsilon \geq 0$. Denoting by d the derivative with respect to \mathbf{u} , the jacobian of \mathbf{F}_ϵ is

$$d\mathbf{F}_\epsilon(\mathbf{u}) = \begin{pmatrix} df_c(u) & -1 & 0 \\ -\epsilon dg(u) & 0 & \epsilon \\ 0 & \epsilon & 0 \end{pmatrix}$$

with characteristic polynomial

$$\det(d\mathbf{F}_\epsilon(\mathbf{u}) - \lambda \mathbb{I}) = -\lambda^3 + df_c(u) \lambda^2 + \epsilon(dg(u) + \epsilon)\lambda - \epsilon^2 df_c(u),$$

where \mathbb{I} denotes the identity matrix. Denoting with λ_1, λ_2 and λ_3 the roots of the polynomial, there hold

$$\begin{aligned}\mathrm{tr} \, d\mathbf{F}_\epsilon(\mathbf{u}) &= df_c(u) = \lambda_1 + \lambda_2 + \lambda_3, \\ \det d\mathbf{F}_\epsilon(\mathbf{u}) &= -\epsilon^2 df_c(u) = \lambda_1 \cdot \lambda_2 \cdot \lambda_3.\end{aligned}$$

Next, we analyze the real part of the eigenvalues λ_1, λ_2 and λ_3 of $d\mathbf{F}_\epsilon(\mathbf{u}_\pm)$. Indeed, from the Lax condition (2.2), it follows

- a. $\mathrm{tr} \, d\mathbf{F}_\epsilon$ computed at \mathbf{u}_+ (respectively \mathbf{u}_-) is negative (resp. positive);
- b. $\det d\mathbf{F}_\epsilon$ computed at \mathbf{u}_+ (resp. \mathbf{u}_-) is positive (resp. negative),

Since the characteristic polynomial has degree 3, there are two cases:

- i. either 3 real roots;
- ii. or 1 real root and 2 complex conjugates.

Next, we limit the investigation to the case \mathbf{u}_+ , the other being analogous.

- i. Thanks to b., we have either 3 positive eigenvalues or 2 negative eigenvalues and 1 positive. The first case is excluded by a.; thus, there are 2 negative and 1 positive roots.
- ii. Let us set $\lambda_1 \in \mathbb{R}$, $\lambda_2 = \alpha + i\beta$ and $\lambda_3 = \alpha - i\beta$ with $\beta > 0$. Then

$$\mathrm{tr} \, d\mathbf{F}_\epsilon = \lambda_1 + 2\alpha < 0 \quad \text{and} \quad \det d\mathbf{F}_\epsilon = \lambda_1(\alpha^2 + \beta^2) > 0.$$

These conditions imply $\lambda_1 > 0$ and $\alpha = \mathrm{Re}\lambda_2 = \mathrm{Re}\lambda_3 < 0$.

Summarizing, at \mathbf{u}_+ , the matrix $d\mathbf{F}_\epsilon$ has a two-dimensional stable manifold and an one-dimensional stable manifold. At \mathbf{u}_- the situation is reversed: the dimension of the stable manifold is 1 and that of the unstable manifold is 2. We infer that the splitting is consistent since the sum of the dimension of the unstable manifold at \mathbf{u}_- and the one of the stable manifold at \mathbf{u}_+ is equal to 4 and the state space has dimension equal to 3.

sec:extfs

2.2. Extended fast system. Next, we extend the fast system (2.5) adding a trivial equation for the parameter ϵ , that is

$$\dot{\mathbf{v}} = \Phi(\mathbf{v}) := (F(\mathbf{u}), \epsilon G(\mathbf{u}), \epsilon H(\mathbf{u}), 0), \quad (2.7)$$

layer

where $\mathbf{v} = (\mathbf{u}, \epsilon) \in \mathbb{R}^3 \times (-\epsilon_0, \epsilon_0)$ and F, G and H are still defined in (2.3). Incidentally, let us observe that system (2.7) is equivalent to (2.5) for $\epsilon > 0$ and to (2.6) for $\epsilon = 0$.

The matrix of the linearization of Φ at \mathbf{v} is

$$d\Phi(\mathbf{v}) = \begin{pmatrix} df_c(u) & -1 & 0 & 0 \\ -\epsilon dg(u) & 0 & \epsilon & w - g(u) \\ 0 & \epsilon & 0 & v \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

with characteristic polynomial at $\mathbf{v}_0 := (\mathbf{u}, 0)$

$$\det(d\Phi(\mathbf{v}_0) - \lambda \mathbb{I}) = \lambda^3(\lambda - df_c(u)).$$

The polynomial has two roots, $\lambda = 0$ and $\lambda = df_c(u)$, which are distinct if $df_c(u) \neq 0$. The multiplicity 3 of the eigenvalue 0 corresponds to the dimension of the slow variable (v, w) plus the dimension of the fictitious variable ϵ ; the multiplicity 1 of the eigenvalue $df_c(u)$, coincides with the dimension of the fast variable u . Moreover, since $\lambda = df_c(u) > 0$ on \mathcal{S}_- and $\lambda = df_c(u) < 0$ on \mathcal{S}_+ , the unstable manifold of \mathbf{u}_- in \mathcal{S}_- and the stable manifold of \mathbf{u}_- in \mathcal{S}_+ have both dimension equal to 1.

For readers' convenience, let us recall a basic definition valid for generic compact manifold M and diffeomorphism f .

`f: normallyhyperbolic`

Definition 2.1. *An f -invariant submanifold Λ of M is said to be **normally hyperbolic** if the restriction $T_\Lambda M$ to Λ of the tangent bundle TM admits a splitting into a sum of three df -invariant sub-bundles,*

$$T_\Lambda M = T\Lambda + E^s + E^u.$$

where $T\Lambda$ is the tangent bundle of Λ and $E^{s,u}$ denotes the stable/unstable bundle, respectively.

Here, we consider compact subsets $\mathcal{K} \Subset \mathcal{S}_0$ that are normally hyperbolic invariant manifolds of the layer problem (2.6).

Since \mathbf{F}_0 restricted to \mathcal{S}_\pm is identically zero, there holds

$$T_{\mathbf{u}}\mathcal{S}_\pm = \ker d\mathbf{F}_0.$$

The subspace $T_{\mathbf{u}}\mathcal{S}_\pm$ is invariant under the action of $d\mathbf{F}_0$ and, therefore, the linear map induced by the linearization

$$Q\mathbf{F}_0(\mathbf{u}) : T_{\mathbf{u}}\mathbb{R}^3/T_{\mathbf{u}}\mathcal{S}_\pm \rightarrow T_{\mathbf{u}}\mathbb{R}^3/T_{\mathbf{u}}\mathcal{S}_\pm$$

is well defined.

The eigenvalues of $d\mathbf{F}_0$ are exactly the nontrivial ones cited before, since

$$d\mathbf{F}_0(\mathbf{u}) = \begin{pmatrix} (df_c \circ h_\pm)(v) & -1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Moreover, a generic element $\mathbf{v} \in T_{\mathbf{u}}\mathcal{S}_\pm$ is $\mathbf{v} = \alpha \bar{\mathbf{u}} + \beta \mathbf{e}_3$ for $(\alpha, \beta) \in \mathbb{R}^2$ where $\bar{\mathbf{u}} := (1, (df_c \circ h_\pm)(v), 0)$.

Next, following [19], we define the *projection map* $\pi^{\mathcal{S}}$ by the splitting

$$T\mathbb{R}^3 = T\mathcal{S}_\pm \oplus N,$$

where N is the complement of $T\mathcal{S}_\pm$ invariant under $d\mathbf{F}_0$. The matrix $d\mathbf{F}_0$ maps the vector $\mathbf{v} = (a, b, c) \in \mathbb{R}^3$ into

$$\begin{pmatrix} (df_c \circ h_\pm)(v) & -1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix} = \begin{pmatrix} (df_c \circ h_\pm)(v)a - b \\ 0 \\ 0 \end{pmatrix}.$$

Therefore, for a nonzero vector (a, b, c) to be invariant under the action of $J\mathbf{F}_0$ it is required $b = c = 0$. Since $(df_c \circ h_{\pm})(v) \neq 0$, the subspace N is thus spanned by $\mathbf{e}_1 := (1, 0, 0)$. Summarizing, the splitting is

$$T\mathbb{R}^3|_{\mathcal{S}_{\pm}} = \langle \bar{\mathbf{u}}, \mathbf{e}_3 \rangle \oplus \langle \mathbf{e}_1 \rangle.$$

Let us denote with \mathcal{C} , \mathcal{C}^s and \mathcal{C}^u and the center, center-stable, center-unstable manifolds generated by the extended fast system Φ . They are locally invariant manifolds containing $\mathcal{K} \times \{0\}$ and tangent to the corresponding center, center-stable, center-unstable eigenspaces of the linearization $d\Phi$ at $\mathbf{v} = (\mathbf{u}, 0) \in \mathcal{K} \times \{0\}$, denoted here by $E_{\mathbf{u}}^c$, $E_{\mathbf{u}}^s \oplus E_{\mathbf{u}}^c$ and $E_{\mathbf{u}}^u \oplus E_{\mathbf{u}}^c$,

In the case under scrutiny, the dimension k^c of the center manifold for the system (2.5) is 0. We recall that $n = 2$ is the dimension of the slow variable (v, w) and $k = k^u + k^s = 1$ equals the dimension of the fast variable. The following statements are true:

1. $\dim \mathcal{C} = n + 1 = 3$;
2. for the surface \mathcal{S}_+ , $k^s = 1$ and $k^u = 0$, so that $\dim \mathcal{C}^s = n + 1 + k^s = 4$ and $\dim \mathcal{C}^u = n + 1 + k^u = 3$;
3. for the surface \mathcal{S}_- , $k^s = 0$ and $k^u = 1$ so that $\dim \mathcal{C}^s = n + 1 + k^s = 3$ and $\dim \mathcal{C}^u = n + 1 + k^u = 4$.

Given constants $p, q \in [1, +\infty)$, let \mathcal{C}^s be a center-stable manifold for Φ in a neighborhood of $\mathcal{K} \times \{0\}$. At this point, we need to introduce also the notion of *family of stable (resp. unstable) manifolds* for \mathcal{C}^s (resp. \mathcal{C}^u).

Definition 2.2. A family $\mathcal{F}^s = \{\Psi^s(\mathbf{v}) : \mathbf{v} \in \mathcal{C}^s\}$ is a C^q -family of C^p -stable manifold for \mathcal{C}^s near \mathcal{K} if

- i. $\Psi^s(\mathbf{v})$ is C^p -manifold for each $\mathbf{v} \in \mathcal{C}^s$;
- ii. $\mathbf{v} \in \Psi^s(\mathbf{v})$ for each $\mathbf{v} \in \mathcal{C}^s$;
- iii. $\Psi^s(\mathbf{v}_1)$ and $\Psi^s(\mathbf{v}_2)$ are disjoint if $\mathbf{v}_1 \neq \mathbf{v}_2$ for each $\mathbf{v}_1, \mathbf{v}_2 \in \mathcal{C}^s$;
- iv. $\Psi^s(\mathbf{u}, 0)$ is tangent to $E_{\mathbf{u}}^s$ at $(\mathbf{u}, 0)$ for each $\mathbf{u} \in \mathcal{S}$;
- v. the set $\{\Psi^s(\mathbf{v}) : \mathbf{v} \in \mathcal{C}^s\}$ is a positively invariant C^q -family of manifolds with respect to the flow Φ .

The case of the unstable manifolds \mathcal{F}^u is defined in an analogous way. These families provide a *foliation* of \mathcal{C}^s and \mathcal{C}^u , i.e.

$$\mathcal{C}^s = \{\Psi^s(\mathbf{v}) : \mathbf{v} \in \mathcal{C}\} \quad \text{and} \quad \mathcal{C}^u = \{\Psi^u(\mathbf{v}) : \mathbf{v} \in \mathcal{C}\}.$$

sec:red0

2.3. Construction of the profile for the reduced system. We want to apply Lemma A.2 in Appendix A to system (2.5) to describe the flow induced on \mathcal{S} when it is given by the graph of a function, i.e. in \mathcal{S}_{\pm} .

Adapting Lemma 5.4 in [5] to (2.5), we infer that the projections π^\pm on the surfaces \mathcal{S}_\pm are given by the multiplication against the matrices

$$\mathbb{A}^\pm = \begin{pmatrix} 0 & dh_\pm(v) & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

so that the *reduced systems* are

$$\mathbf{u}' = \mathbb{A}^\pm \begin{pmatrix} 0 \\ G \\ H \end{pmatrix} = \begin{pmatrix} dh_\pm(v) G(h_\pm(v), v, w) \\ G(h_\pm(v), v, w) \\ H(h_\pm(v), v, w) \end{pmatrix}$$

In components, for any $(u, v, w) \in \mathcal{S}_\pm$, the above systems become

$$\begin{cases} u' = \frac{w - (g \circ h_\pm)(v)}{(df_c \circ h_\pm)(v)} \\ v' = w - (g \circ h_\pm)(v) \\ w' = v. \end{cases} \quad (2.8) \quad \boxed{\text{reduced}}$$

In what follows, we refer to the vector field as \mathbf{F}_R .

Remark 2.3. In the case under investigation, the computations yielding to (2.8) reduces to differentiate (2.4)₁ along the profile lying on \mathcal{S}_\pm . Hence, for $u = h_\pm(v)$, and using (2.4)₂ we end up with

$$u' = \frac{w - (g \circ h_\pm)(v)}{(df_c \circ h_\pm)(v)}$$

and thus (2.8). On the other hand, Lemma A.2 in Appendix A refers to a general framework, where the vector field in (2.5) merely depends in a regular way on ϵ .

Upon observation, it is readily seen that the critical points of the *limiting slow system* (2.4), or equivalently (2.8), where we have desingularized the problem, are

$$\mathbf{u}_\pm := (u_\pm, 0, g(u_\pm)) \in \mathcal{S}_\pm,$$

and hence they coincide with the ones of the perturbed slow systems (2.3). Theorem A.3 in Appendix guarantees the existence for $\epsilon > 0$ sufficiently small of a sequence of critical points \mathbf{u}_ϵ for the perturbed vector field in (2.3), smoothly depending on ϵ , and which reduces for $\epsilon = 0$ to the critical points of (2.4). In our case, again in view of the independence from ϵ of the vector field in (2.3), we have $\mathbf{u}_\pm = \mathbf{u} = \mathbf{u}_\epsilon$, without invoking Theorem A.3. However, in order to define the local stable/unstable manifolds of \mathbf{u}_ϵ needed for the connection, we shall refer to the general framework of Theorem A.4 and therefore one has to check that $\lambda = 1$ is not an eigenvalue of the Jacobian

$d\mathbf{F}_R(\mathbf{u}_\pm)$. To this end, we compute the Jacobian of the vector fields in (2.8)

$$d\mathbf{F}_R(\mathbf{u}_\pm) = \begin{pmatrix} 0 & -dg/df_c^2 & 1/df_c \\ 0 & -dg/df_c^2 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

where df_c and dg are computed at u_\pm . The corresponding characteristic polynomials p_\pm are

$$p_\pm(\lambda) := -\lambda q_\pm(\lambda) \quad \text{with} \quad q_\pm(\lambda) = \lambda^2 + \frac{dg(u_\pm)}{df_c(u_\pm)} \lambda - 1.$$

These have three roots: the trivial one, given by $\lambda_\pm^0 = 0$, and the roots of q_\pm , which can be expressed as

$$\begin{aligned} \lambda_\pm^1 &= -\frac{\sqrt{dg^2 + 4df_c^2} + (\operatorname{sgn} df_c) dg}{2|df_c|}, \\ \lambda_\pm^2 &= \frac{\sqrt{dg^2 + 4df_c^2} - (\operatorname{sgn} df_c) dg}{2|df_c|}, \end{aligned}$$

where df_c and dg are computed at u_\pm . Recalling the Lax condition (2.2) and the monotonicity assumption (1.4) on function g , a direct computation shows that

$$\lambda_-^1 < -1 < \lambda_+^1 < 0 < \lambda_-^2 < 1 < \lambda_+^2,$$

so that, in particular, $\lambda_\pm^2 \neq 1$.

Therefore we can apply Theorem A.4 to characterize the local stable-unstable manifolds of the perturbed critical points. These manifolds are of dimension 1 for both \mathbf{u}_\pm and thus the critical points for the perturbed system are saddles in the slow directions.

Now we pass to the study of the reduced system (2.8) to briefly recast the results by Lattanzio et al. [11, 12] in the present framework (see also [8]). To this end, we focus on the case of sufficiently large shocks so that the profile exhibits a sub-shock for the variable u , case for which GSPT can be directly applied, because the two branches of the profile belong to \mathcal{S}_- and \mathcal{S}_+ , respectively.

In the case of the *singularly perturbed systems* (2.3), we prove the existence of the heteroclinic orbit as a perturbation of the so-called *singular heteroclinic orbit*. The latter consists of orbits of the reduced systems lying on the two surfaces \mathcal{S}_\pm of the critical manifold \mathcal{S} , linked by a heteroclinic connection of the *layer system* (2.6).

For completeness, we adapt here the results proved in [11] for system (1.3) with $g(u) = u$ and in [12] for general increasing functions g . For readers' convenience, we observe that the variable (u, v, w) of the present paper corresponds to $(u, -q, z)$ of [11, 12].

Let (v_{\pm}, w_{\pm}) be the maximal solutions to the problems

$$v' = w - (g \circ h_{\pm})(v), \quad w' = v, \quad (2.9) \quad \boxed{\text{wpm}}$$

with the asymptotic conditions $(v_{\pm}, w_{\pm})(\pm\infty) = (0, g(u_{\pm}))$, describing the dynamics of the reduced system (2.8) along the surfaces \mathcal{S}_{\pm} .

Functions v_{-} , w_{\pm} are decreasing and v_{+} is increasing. In addition, there exist $x_{\pm} \in \mathbb{R}$ such that

$$w_{\pm}(x_{\pm}) - v'_{\pm}(x_{\pm}) = g(u_{\pm}) \quad \text{and} \quad v_{\pm}(x_{\pm}) = -f_c(u_{\pm})$$

In particular, the solution (v_{-}, w_{-}) is defined in $(-\infty, x_{*}]$ and the solution (v_{+}, w_{+}) is defined in $[x_{*}, +\infty)$.

As a consequence of the monotonicity of functions v_{\pm} , there holds

$$w_{-}(x_{*}) \leq g(u_{*}) \leq w_{+}(x_{*}).$$

The complete heteroclinic orbit is built by matching two branches of the maximal solutions (v_{\pm}, w_{\pm}) at a given point x_{*} which can be arbitrarily chosen taking advantage of translation invariance.

teo3.5

Proposition 2.4. *Assume hypotheses (1.4). Then, there exists a function (v, w) with $v \in C^0(\mathbb{R}) \cap C^1(\mathbb{R} \setminus \{x_{*}\})$ and $w \in C^1(\mathbb{R}) \cap C^2(\mathbb{R} \setminus \{x_{*}\})$ solving the first order system (not in normal form)*

$$(f_c \circ g^{-1})(w - v') = v, \quad w' = v,$$

satisfying the asymptotic conditions $(v, w)(\pm\infty) = (0, g(u_{\pm}))$. Moreover, the function $v' = w''$ has at most a jump discontinuity at x_{} .*

The existence theorem of propagating fronts for system (1.3) with $\epsilon = 0$ is a simple consequence of the above construction.

teo:existsinglim

Theorem 2.5. *Assume hypotheses (1.4). For any u_{\pm} satisfying the Lax condition (1.7), there exists a propagating front for system (1.3) with $\epsilon = 0$. The profile (u, v) is unique up to translation and the speed c is given by the Rankine-Hugoniot condition (1.6).*

Moreover, the component u belongs to $C^1(\mathbb{R} \setminus \{x_{}\})$, eventually with a jump discontinuity at x_{*} .*

For the complete proofs of Proposition 2.4 and Theorem 2.5, we refer to [11, 12]. Here, we limit to sketch the basic steps in the matching procedure with the aim of providing support to the subsequent parts.

We define a profile w by matching together the two maximal solutions

$$w(x) = \begin{cases} w_{-}(x) & x < x_{*} \\ w_{+}(x) & x \geq x_{*}, \end{cases}$$

relying on the monotonicity of maximal solutions w_{\pm} , such that $w(x_*) = w_*$. The latter equality can be written also as

$$g(u_r) + v'_r = g(u_\ell) + v'_\ell.$$

Analogously, the continuity of w' at x_* gives

$$w'_+(x_*) = v_r = v_\ell = w'_-(x_*).$$

Since we are investigating a discontinuous profile u , we know that

$$v_* := v_\ell = v_r < 0, \quad (2.10) \quad \text{vastnegative}$$

and

$$u_r = g^{-1}(w_* - v'_+(x_*)) < u_* < g^{-1}(w_* - v'_-(x_*)) = u_\ell. \quad (2.11) \quad \text{eq:ulurdiff}$$

Therefore, there exists $\delta > 0$ sufficiently small such that, along the constructed profile, we have

$$h_+(v) \in [u_+, u_* - \delta], \quad h_-(v) \in [u_* + \delta, u_-],$$

and GSPT is applicable.

Remark 2.6. The present analysis is valid also for the particular case of the Hamer model with linear coupling (1.2), namely for $f(u) = u^2/2$, $g(u) = u$. In that case, the slow manifolds are given by:

$$\mathcal{S}_{\pm} = \{(u, v, w) \in \mathbb{R}^3 \mid u = \mp \sqrt{1 + 2v}\}.$$

If we choose the asymptotic states as $u_{\pm} = (\mp 1, 0, \mp 1)$, we obtain $u^* = 0$ and $c = 0$. In [8, 11], the authors showed that the profile increases its regularity as the strength of the shock decreases. Moreover, for that specific model, from [8] we know that u has precisely an ammissible jump discontinuity at a single point if $|u_+ - u_-| > \sqrt{2}$, which is the case we are referring to here. Therefore we get only the C^1 -continuity of w . The maximal solution w_+ decreases from zero toward -1 at $+\infty$ and w_- decreases from 1 at $-\infty$ toward 0. Due to the particular choice of the flux f , and thus of the inverse function $h_{\pm}(v)$, we get $u_\ell = -u_r$. We shall give a partial answer in the case of small shocks in Section 3, when we shall discuss the existence of the profile for sufficiently small shocks and arbitrary diffusion.

subsec:singular

2.4. Construction and persistence of the orbit. In the previous section we have constructed the first two pieces of the singular heteroclinic orbit, namely orbits of the reduced systems \mathbf{F}_R lying on \mathcal{S}_{\pm} . Here, we aim to substitute the instantaneous jump for u among these branches with an appropriate smoothed version described by the limiting equation (2.4) of the original differential equations (2.3) in the regime $\epsilon \rightarrow 0^+$.

The limiting fast system (2.6) has critical points given by $(h_{\pm}(v), v, w)$ for arbitrary v, w . Therefore, we make use of it to connect the two manifolds \mathcal{S}_{\pm} at points $\mathbf{u}_{\ell} = (u_{\ell}, v_*, w_*) \in \mathcal{S}_{-}$ and $\mathbf{u}_r = (u_r, v_*, w_*) \in \mathcal{S}_{+}$, where $u_{\ell} = h_{-}(v_*)$ and $u_r = h_{+}(v_*)$. The dynamics are completely described by the reduced equation

$$\dot{u} = F(u, v_*, w) = f_c(u) - f_c(u_{\pm}) - v_*. \quad (2.12) \quad \text{eq:ODE1non}$$

Thus, for fixed $v_* > f_c(u_*) - f_c(u_-)$, it is easily checked that

$$\begin{aligned} F(u, v_*, w) &= 0 \quad \text{if and only if} \quad u = u_{\ell} \quad \text{or} \quad u = u_r, \\ F(u, v_*, w) &< 0 \quad \text{for} \quad u \in (u_r, u_{\ell}), \end{aligned}$$

with the opposite sign for $u \in (-\infty, u_r) \cup (u_{\ell}, \infty)$. We infer the existence of a global and decreasing solution u_0 to (2.12) verifying the asymptotic conditions $u_0(-\infty) = u_{\ell}$ and $u_0(+\infty) = u_r$.

rem:Hamer_u0

Example 2.7 (Hamer model). Let $u_{\mp} = \pm 1$. As a consequence of the Rankine–Hugoniot (1.6), the speed c is zero and

$$f_0(u) = f(u) - 0 \cdot u = \frac{1}{2}u^2.$$

Moreover, since $|u_- - u_+| = 2 > \sqrt{2}$, the inviscid profile undergoes a Lax sub-shock between u_{ℓ} and $u_r = -u_{\ell} < 0$. For any fixed value $v_* \in (-\frac{1}{2}, 0)$, we can explicitly solve the equation (2.12) for u . Indeed, setting $v_* := -\frac{1}{2}(u_{\ell}^2 - 1)$ for some $u_{\ell} \in (0, 1)$, there holds

$$\dot{u} = \frac{1}{2}(u^2 - 1) - v_* = \frac{1}{2}(u^2 - u_{\ell}^2).$$

Separating the variables and integrating by parts, we obtain

$$u_0(x) = -u_{\ell} \cdot \frac{e^{u_{\ell}x} - 1}{e^{u_{\ell}x} + 1} = -u_{\ell} \tanh\left(\frac{1}{2}u_{\ell}x\right),$$

which satisfies the required asymptotic conditions $u_0(\pm\infty) = \mp u_{\ell}$.

Let $a, b \in \mathbb{R}$ with $a \leq b$. Then, given $\delta \in (0, f_c(u_{\pm}))$, the compact normally hyperbolic invariant manifolds \mathcal{K}_{\pm} have the form

$$\mathcal{K}_{\pm} = \{\mathbf{u} \in \mathcal{S}_{\pm} : (v, w) \in [\delta - f_c(u_{\pm}), 0] \times [a, b]\}.$$

It is clearly seen that the two parameter sets coincide.

Let $\mathcal{W}_{-,0}^u$ be the unstable manifold of \mathbf{u}_{-} and $\mathcal{W}_{+,0}^s$ the stable manifold of \mathbf{u}_{+} with respect to the reduced slow system (2.8). Then, the *singular unstable manifold* of \mathbf{u}_{-} and *singular stable manifold* of \mathbf{u}_{+} are given by

$$N_{-,0}^u = \bigcup_{\mathbf{u} \in \mathcal{W}_{-,0}^u} \Psi^u(\mathbf{u}) \quad \text{and} \quad N_{+,0}^s = \bigcup_{\mathbf{u} \in \mathcal{W}_{+,0}^s} \Psi^s(\mathbf{u}),$$

where $\Psi^{u,s}(\mathbf{u})$ are the unstable/stable fibers of the limiting fast system (2.6) based at \mathbf{u} , which are tangent to the eigenvectors associated to the non-trivial eigenvalue of the Jacobian $d\mathbf{F}_0(\mathbf{u})$ with an appropriate sign choice.

The construction of the singular heteroclinic orbit is complete if and only if we show that the intersection of the manifolds $N_{-,0}^u$ and $N_{+,0}^s$ is transversal along the solution of (2.12), namely upon construction of the smooth connection between u_ℓ and u_r recalled above.

For $\mathbf{u} \in \mathcal{S}_*$, the non trivial eigenvalue of the Jacobian $d\mathbf{F}_0(\mathbf{u})$ is $\lambda = df_c(u)$ which is negative in \mathcal{S}_+ and positive in \mathcal{S}_- . At $\mathbf{u} \in \mathcal{S}_-$, the eigenvector relative to the eigenvalue $df_c(u)$ is readily seen to be $\mathbf{e}_1 = (1, 0, 0)$, so that the unstable fiber $\Psi^u(\mathbf{u})$ is explicitly given by $\mathbf{u} + t\mathbf{e}_1$ for $t \in \mathbb{R}$. The same computation is valid for $\mathbf{u} \in \mathcal{S}_+$ so that the elements of $\Psi^s(\mathbf{u})$ have the form $\mathbf{u} + t\mathbf{e}_1$ for some $t \in \mathbb{R}$.

Example 2.8 (Hamer model). For the Hamer model \mathcal{K}^\pm are of the form

$$\mathcal{K}_\pm = \left\{ \mathbf{u} \in \mathcal{S}_\pm : (v, w) \in \left[\delta - \frac{1}{2}, 0 \right] \times [a, b] \right\}.$$

for $\delta \in (0, \frac{1}{2})$. The unstable/stable fibers $\Psi^{u,s}(\mathbf{u})$ at $u_\pm = (\mp 1, 0, \mp 1)$ are also tangent to \mathbf{e}_1 .

In view of the previous discussion, for $v_* > f_c(u_*) - f_c(u_-)$, there exists a connection u_0 , solution to (2.12), between the points

$$\mathbf{u}_\ell = (u_\ell, v_*, w_*) \in \mathcal{W}_{-,0}^u \subset \mathcal{K}_- \quad \text{and} \quad \mathbf{u}_r = (u_r, v_*, w_*) \in \mathcal{W}_{+,0}^s \subset \mathcal{K}_+,$$

where $u_\ell = h_-(v_*)$ and $u_r = h_+(v_*)$. Such connections are uniquely determined in view of the monotonicity of the component w along the stable/unstable manifolds $\mathcal{W}_{+,0}^s$ and $\mathcal{W}_{-,0}^u$.

The subsequent Theorem –an ultra-simplified version of [19, Theorem 4.1] which for completeness we stated in the Appendix (see Theorem A.6)– gives conditions to prove transversality in the regime $\epsilon = 0$.

thm:trans

Theorem 2.9. *Let $\phi(\mathcal{W}_{-,0}^u)$ and $\phi(\mathcal{W}_{+,0}^s)$ denote the (v, w) -coordinates of the manifolds $\mathcal{W}_{-,0}^u \subset \mathcal{S}_-$ and $\mathcal{W}_{+,0}^s \subset \mathcal{S}_+$, respectively. Then, the manifolds $N_{-,0}^u$ and $N_{+,0}^s$ intersect transversally at the points of the heteroclinic orbit if and only if*

$$T_{(v_*, w_*)}\phi(\mathcal{W}_{-,0}^u) \cap T_{(v_*, w_*)}\phi(\mathcal{W}_{+,0}^s) = \{0\}. \quad (2.13) \quad \text{nointer}$$

For the proof, see [19]. In the sequel, we check the validity of (2.13).

Remark 2.10. The intersection of $N_{-,0}^u$ and $N_{+,0}^s$ at a point of the form $\mathbf{u} = (u_0(\bar{x}), v_*, w_*)$ (for some generic $\bar{x} \in \mathbb{R}$) is transversal if and only if $T_{\mathbf{u}}N_{-,0}^u + T_{\mathbf{u}}N_{+,0}^s = \mathbb{R}^3$, that is if and only if

$$\dim(T_{\mathbf{u}}N_{-,0}^u + T_{\mathbf{u}}N_{+,0}^s) = 3.$$

Following the notation in [19], the equality

$$\begin{aligned} \dim(T_{\mathbf{u}}N_{-,0}^u + T_{\mathbf{u}}N_{+,0}^s) &= \dim(T_{\mathbf{u}}N_{-,0}^u) + \dim(T_{\mathbf{u}}N_{+,0}^s) \\ &\quad - \dim(T_{\mathbf{u}}N_{-,0}^u \cap T_{\mathbf{u}}N_{+,0}^s) \end{aligned}$$

implies that the intersection in (2.13) is transversal if and only if

$$\dim(T_{\mathbf{u}}N_{-,0}^u \cap T_{\mathbf{u}}N_{+,0}^s) = d,$$

where d in our case is given by

$$d = j_-^u + k_-^u + j_+^s + k_+^s - n - k = 1 + 1 + 1 + 1 - 2 - 1 = 1,$$

with j_-^u , k_-^u , j_+^s and k_+^s describing the dimensions of the stable and unstable reduced manifolds of the singular points \mathbf{u}_{\pm} . In the general setting, Theorem A.6 guarantees the needed transversal intersection along points of the heteroclinic orbit if and only if there exist exactly $d - 1$ linearly independent solutions $\xi \in T_{(v_*, w_*)}\phi(W_{-,0}^u) \cap T_{(v_*, w_*)}\phi(W_{+,0}^s)$ of the equation

$$(M, \xi) = 0,$$

where $M \in \mathbb{R}^2$ is defined by

$$M := \int_{\mathbb{R}} \psi(\xi) \partial_{v,w} \{f_c(u) - f_c(u_-) + v\} d\xi$$

and ψ is defined as the unique bounded solution of

$$\psi' = -df_c(u_0(x)) \psi.$$

Here $d - 1 = 0$ and $\partial_{v,w}(f_c(u) - f_c(u_-) + v) = (1, 0)$. Moreover

$$\psi(x) = \psi(0) \exp \left\{ - \int_0^x df_c(u_0) d\xi \right\}.$$

Since $u_r < u_0(x) < u_\ell$ and df_c is monotone, then

$$df_c(u_r) < df_c(u_0(x)) < df_c(u_\ell),$$

which implies

$$\psi(0)e^{-df_c(u_\ell)x} < \psi(0)e^{-\int_0^x df_c(u_0)ds} < \psi(0)e^{-df_c(u_r)x}.$$

In view of the conditions $df_c(u_r) < 0 < df_c(u_\ell)$, we can conclude that the only ψ which is globally bounded in \mathbb{R} is the trivial one, that is $\psi \equiv 0$ and as a consequence the vector M is identically 0. In other words, in the present case, the requirements of Theorem A.6 reduce to (2.13).

Analyzing the intersection in (2.13), this condition is equivalent to inquire the existence of a couple $(v_0, w_0) \in R := [\delta - f_c(u_{\pm}), 0] \times [a, b] \subset \mathbb{R}^2$ such that the points $(h_-(v), v, w) \in \mathcal{W}_{-,0}^u$ and $(h_+(v), v, w) \in \mathcal{W}_{+,0}^s$ are connected by a heteroclinic orbit of the layer problem (2.6), i.e. a solution u_0 to (2.12) for some constant $w \in \mathbb{R}$. By construction, both the maximal solution w_- (describing the unstable manifold $\mathcal{W}_{-,0}^u$) and the maximal solution w_+

(describing the stable manifold $\mathcal{W}_{+,0}^s$) are monotone decreasing. Hence, since w_- decreases from $g(u_-)$ at $-\infty$ toward w_* and w_+ decreases from w_* toward $g(u_+)$ at $+\infty$, we conclude that the only point in R verifying the above conditions is

$$(v_0, w_0) = (v_r, w_*) = (v_\ell, w_*) = (v_*, w_*).$$

In other words, in order to construct the required smooth connection we are looking for, we have obtained the same condition which locates the discontinuity in u of the profile for $\epsilon = 0$.

Finally, computing the vector field $(G, H) = (w - g(u), v)$ at two given points $(u_\ell, v_*, w_*) \in \mathcal{W}_{-,0}^u$ and $(u_r, v_*, w_*) \in \mathcal{W}_{+,0}^s$ and taking a linear combination of them, we infer

$$\begin{aligned} & \alpha(G, H)(u_\ell, v_*, w_*) + \beta(G, H)(u_r, v_*, w_*) \\ &= ((\alpha + \beta)w_* + \alpha g(u_\ell) + \beta g(u_r), (\alpha + \beta)v_*). \end{aligned}$$

Since $v_* < 0$, see (2.10), $u_r < u_\ell$, see (2.11), and g is monotone increasing, see (1.4), the two vectors are linearly independent which implies (2.13) and then, invoking Theorem 2.9, $N_{-,0}^u$ and $N_{+,0}^s$ intersect transversally. Finally, the heteroclinic orbit for the reduced system at $\epsilon = 0$ persists for $\epsilon > 0$ sufficiently small and the proof of Theorem 1.2 is complete. Indeed, thanks to GSPT, we obtain the existence of the perturbed manifolds $N_{-, \epsilon}^u$ and $N_{+, \epsilon}^s$, and their transversality along a transversal heteroclinic orbit is given by Theorem A.5 (details on the construction of the manifolds $N_{-, \epsilon}^u$ and $N_{+, \epsilon}^s$ can be found in [19]).

Example 2.11 (Hamer model). The same conclusion holds for the Hamer model, for which $R = [\delta - \frac{1}{2}, 0] \times [a, b]$ and $d = 1$. Thus we refer to Theorem 2.8 to check the transversality of $N_{+,0}^s$ and $N_{-,0}^u$. As before, the condition of Theorem 2.8 is equivalent to find $(v_0, w_0) \in R$ such that points $(\sqrt{1+2v_0}, v_0, w_0) \in \mathcal{W}_{-,0}^u$ and $(-\sqrt{1+2v_0}, v_0, w_0) \in \mathcal{W}_{+,0}^s$ are connected by

$$u_0(x) = -u_\ell \tanh\left(\frac{1}{2}u_\ell x\right).$$

As said before, the two maximal solution of the reduced systems w_\pm are monotone decreasing, namely w_+ decreases from zero toward -1 at $+\infty$ and w_- decreases from 1 at $-\infty$ toward 0. Hence, we can conclude that the only point in R verifying the conditions of Theorem 2.8 is $(v_0, w_0) = (v^*, 0)$. The vector field $(G, H) = (w - u, v)$ computed at $(u_\ell, v^*, 0) \in \mathcal{W}_{-,0}^u$ is given by $(-u_\ell, v^*)$, and at $(u_r, v^*, 0) \in \mathcal{W}_{+,0}^s$ is given by $(-u_r, v^*) = (u_\ell, v^*)$. Since $v^* \in [\delta - \frac{1}{2}, 0]$ is not zero, the two vectors are linearly independent and this gives (2.13). We stress that $w = 0$, guaranteeing the existence of the smooth

connection between u_ℓ and u_r , is the same condition found in [8] to locate the admissible sub-shock for the linear, inviscid Hamer model.

sec:smallshocks

3. VISCOUS RADIATING PROFILE FOR SMALL SHOCKS

In the previous sections we proved the existence of viscous radiating profiles for sufficiently small viscosity and large shocks, in particular under the hypothesis that the profiles in the variable u has a sub-shock. Here we complement such result by proving this existence in the case of $O(1)$ -viscosity, fixed to be equal to 1, and sufficiently small shocks via a bifurcation argument with respect to the strength of the shock. This leads to the following system:

$$\begin{cases} \partial_t u + \partial_x f(u) - \partial_{xx} v = \partial_{xx} u \\ v - \partial_{xx} v = \partial_x g(u), \end{cases} \quad (3.1) \quad \text{PDE2}$$

where we assume always hypotheses (1.4). We shall prove the existence of solutions to (3.1) in form of travelling waves, i.e. $(u, v)(x, t) = (u(\xi), v(\xi))$ where $\xi := x - ct$ (with a slight abuse of notation), when the strength of the underlying Lax shock is sufficiently small:

$$\begin{aligned} u(\pm\infty) &= u_\pm, \quad v(\pm\infty) = 0, \\ c &= \frac{f(u_+) - f(u_-)}{u_+ - u_-}, \\ f'(u_+) &< c < f'(u_-) \iff u_+ < u_-, \\ \delta &:= u_+ - u_- < 0, \end{aligned}$$

δ being the (negative) bifurcation parameter. In what follows, for the sake of simplicity, we shall focus on the Hamer model, namely when $f(u) = u^2/2$ and $g(u) = u$, being the general case presented above analogous.

System (3.1) reduces to (1.2):

$$\begin{cases} \partial_t u + \partial_x \left(\frac{1}{2} u^2 \right) - \partial_{xx} v = \partial_{xx} u \\ v - \partial_{xx} v = u, \end{cases} \quad (3.2) \quad \text{PDE22}$$

and the Rankine-Hugoniot condition becomes:

$$c = \frac{1}{2} \frac{(u_+)^2 - (u_-)^2}{u_+ - u_-} = \frac{1}{2} (u_+ + u_-). \quad (3.3) \quad \text{c}$$

Moreover, the Lax condition rewrites:

$$u_+ < c < u_-. \quad (3.4) \quad \text{lax3}$$

In particular, we prove a simplified version of Theorem 1.3, that is:

thm:main2

Theorem 3.1. *Assume (3.3), (3.4). Moreover, let u_- and u_+ be such that $|u_+ - u_-|$ is sufficiently small. Then the parabolic-elliptic system (3.2) supports propagating fronts with speed c given by the Rankine-Hugoniot relation (3.3).*

The remaining part of this section is devoted to the proof of this result.

To this aim, let us start by recalling the dynamical system solved by the profile:

$$\begin{cases} -cu' + \left(\frac{1}{2}u^2\right)' - v' = u'' \\ -v'' + v = u'. \end{cases}$$

Integrating the first equation in $d\xi$, $\xi = x - ct$, from $\pm\infty$ we get:

$$\begin{cases} u' = \frac{1}{2}(u^2 - u_{\pm}) - v - c(u - u_{\pm}) \\ -v'' + v = u'. \end{cases} \quad (3.5) \quad \boxed{\text{ode}}$$

Moreover, we can express the speed of the wave c in (3.3) as a function of u_+ and δ as follows

$$c = \frac{1}{2}(2u_+ - \delta)$$

and therefore

$$\begin{cases} u' = \frac{1}{2}(u - u_+)[(u - u_+) + \delta] - v \\ -v'' + v = u'. \end{cases}$$

With the notation $\tilde{u} = u - u_+$ and $z = \tilde{u} + v'$, the dynamical system becomes:

$$X' = F(X; \delta), \quad (3.6) \quad \boxed{\text{ODE}}$$

where $X = (z, v, \tilde{u})$ and

$$F(z, v, \tilde{u}; \delta) = \left(v, z - \tilde{u}, \frac{1}{2}\tilde{u}^2 + \frac{1}{2}\tilde{u}\delta - v\right).$$

Hence we observe that $p_1 = (0, 0, 0)$ is a critical point for any δ , which corresponds to the point $(u_+, 0, u_+)$ in the original variables. Moreover, depending on δ , we have two different situations:

- 1) if $\delta = 0$ then p_1 is the only (trivial) critical point;
- 2) if $\delta < 0$, then a second critical point $p_2 = (-\delta, 0, -\delta)$ bifurcates from the trivial one, the latter corresponds to the point $(u_-, 0, u_-)$ in the original variables.

We want to transform (3.6) into its normal form performing a center manifold reduction, and prove a transcritical bifurcation that occurs at $\delta = 0$. This will then imply the existence of the desired heteroclinic orbit, connecting p_2 at $-\infty$ to p_1 at $+\infty$. To this end we start by rewriting (3.6) with respect to the eigenbasis of the linearized system at the trivial critical point for $\delta = 0$.

Let $A = A(z, v, \tilde{u}; \delta)$ be the Jacobian of F at (z, v, \tilde{u}) , namely

$$A(z, v, \tilde{u}; \delta) = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & -1 \\ 0 & -1 & \tilde{u} + \frac{1}{2}\delta \end{pmatrix}$$

The characteristic equation at $p_1 = (0, 0, 0)$ is given by

$$-\lambda^3 + \frac{1}{2}\delta\lambda^2 + 2\lambda - \frac{1}{2}\delta = 0.$$

We study the real part of the eigenvalues at p_1 using the information coming from $\text{tr}(A|_{p_1})$ and $\det(A|_{p_1})$. Since $\text{tr}(A|_{p_1}) = \lambda_1 + \lambda_2 + \lambda_3 = \frac{1}{2}\delta < 0$ and $\det(A|_{p_1}) = \lambda_1\lambda_2\lambda_3 = -\frac{1}{2}\delta > 0$, we have two possibilities:

- 1) one real positive eigenvalue and two complex and conjugates eigenvalues with negative real part;
- 2) two real negative eigenvalues and one real positive eigenvalue.

Therefore at p_1 we have two stable direction and one unstable direction. For $p_2 = (-\delta, 0, -\delta)$ the situation is clearly reversed, being $\text{tr}(A|_{p_2}) = -\frac{1}{2}\delta > 0$ and $\det(A|_{p_2}) = \frac{1}{2}\delta < 0$. Hence at p_2 we have two unstable direction and one stable direction.

For $\delta = 0$ the Jacobian at p_1 reduces to

$$A(0, 0, 0; 0) = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}$$

with eigenvalues $\lambda_1 = 0, \lambda_2 = \sqrt{2}, \lambda_3 = -\sqrt{2}$ and corresponding eigenvectors given by $(1, 0, 1)$, $(-1, \sqrt{2}, 1)$ and $(1, \sqrt{2}, -1)$. Therefore, setting $Y := (w_1, w_2, w_3)$, the desired change of basis is defined by the explicit matrix

$$C = \begin{pmatrix} 1 & -1 & 1 \\ 0 & \sqrt{2} & \sqrt{2} \\ 1 & 1 & -1 \end{pmatrix}$$

giving raise to

$$Y = C^{-1}X. \tag{3.7} \quad \boxed{\text{coo}}$$

As a consequence, our original system (3.6) becomes

$$Y' = BY + C^{-1}\tilde{F}(CY; \delta),$$

where $B := C^{-1}AC$ and $\tilde{F}(X; \delta) = F(X; \delta) - A(0, 0, 0) \cdot X$, namely:

$$\begin{aligned} \frac{d}{dx} \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & -\sqrt{2} & 0 \\ 0 & 0 & +\sqrt{2} \end{pmatrix} \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} \\ &+ \frac{1}{4} \begin{pmatrix} (w_1 + w_2 - w_3)^2 + \delta(w_1 + w_2 - w_3) \\ \frac{1}{2}(w_1 + w_2 - w_3)^2 + \frac{1}{2}\delta(w_1 + w_2 - w_3) \\ -\frac{1}{2}(w_1 + w_2 - w_3)^2 - \frac{1}{2}\delta(w_1 + w_2 - w_3) \end{pmatrix} \end{aligned} \tag{3.8} \quad \boxed{\text{DEC}}$$

In this way the system consists of a linear part and a perturbation. The *Center Manifold Theorem* guarantees the existence for δ sufficiently small of two C^1 -functions $\psi_2(w_1, \delta)$ and $\psi_3(w_1, \delta)$ such that

$$w_2 = \psi_2(w_1, \delta), \quad w_3 = \psi_3(w_1, \delta),$$

and the following tangency conditions hold:

$$\begin{aligned} \psi_2(0, 0) &= \psi_3(0, 0) = 0; \\ \frac{\partial \psi_2}{\partial w_1}(0, 0) &= \frac{\partial \psi_3}{\partial w_1}(0, 0) = 0; \\ \frac{\partial \psi_2}{\partial \delta}(0, 0) &= \frac{\partial \psi_3}{\partial \delta}(0, 0) = 0. \end{aligned} \tag{3.9} \quad \boxed{\text{eq:tang}}$$

Thus for $\delta < 0$ small enough, ψ_2 and ψ_3 are tangent in $(0, 0)$ to the plane identified by $(w_1, \delta) = (0, 0)$. Moreover, the Center Manifold Theorem shows that, again for $\delta < 0$ small, in a neighborhood of the non hyperbolic critical point $(0, 0, 0)$, the original system (3.6), rewritten as in (3.8), is C^1 topologically conjugate to the following decoupled system:

$$\begin{cases} w'_1 = \frac{1}{4} \{w_1 + \psi_2(w_1, \delta) - \psi_3(w_1, \delta)\}^2 + \frac{1}{4} \delta \{w_1 + \psi_2(w_1, \delta) - \psi_3(w_1, \delta)\} \\ w'_2 = -\sqrt{2} w_2, \\ w'_3 = +\sqrt{2} w_3. \end{cases} \tag{3.10} \quad \boxed{\text{G}}$$

Thus, to study the qualitative behaviour of the flow given by (3.6) in a neighborhood of the non hyperbolic critical point $(0, 0, 0)$ and $\delta < 0$ sufficiently small, we are reduced to the flow on the center manifold, which is given by (3.10)₁. We want to apply *Sotomayor Theorem* [17, p. 338] to see which type of bifurcation occurs for $\delta = 0$. For completeness we report the statement below.

Theorem 3.2 (Sotomayor). *Suppose that $F(X_0, \delta_0) = 0$ and that the 3×3 matrix $A = JF(X_0, \delta_0)$ has a simple eigenvalue $\lambda = 0$ with eigenvector v and that A^T has an eigenvector w corresponding to the eigenvalue $\lambda = 0$. Furthermore, suppose that A has one eigenvalue with positive real part and one with negative real part and that the following conditions are satisfied:*

$$\begin{aligned} w^T \cdot F_\delta(X_0, \delta_0) &= 0; \\ w^T \cdot [DF_\delta(X_0, \delta_0)v] &\neq 0; \\ w^T \cdot [D^2F(X_0, \delta_0)](v, v) &\neq 0. \end{aligned} \tag{3.11} \quad \boxed{\text{W}}$$

Then there is a smooth curve of equilibrium points of $X' = F(X; \delta)$ in $\mathbb{R}^3 \times \mathbb{R}$ passing through (X_0, δ_0) and tangent to the hyperplane $\mathbb{R}^3 \times \{\delta_0\}$. Depending on the sign of the expressions in (3.11), there are no equilibrium points near X_0 when $\delta < \delta_0$ (or when $\delta > \delta_0$) and there are two equilibrium points near X_0 when $\delta > \delta_0$ (or when $\delta < \delta_0$). The two equilibrium points of the vector field $F(X; \delta)$ are hyperbolic and have stable manifolds of dimension one and two respectively; i.e., the system $X' = F(X; \delta)$ experiences a transcritical bifurcation at the equilibrium point X_0 as the parameter δ passes through the bifurcation value $\delta = \delta_0$.

Introducing the same notation of the theorem, with $\hat{F}(w_1, w_2, w_3; \delta)$ denoting the vector field

$$\begin{pmatrix} \frac{1}{4} \{w_1 + \psi_2(w_1, \delta) - \psi_3(w_1, \delta)\}^2 + \frac{1}{4} \delta \{w_1 + \psi_2(w_1, \delta) - \psi_3(w_1, \delta)\} \\ -\sqrt{2} w_2 \\ +\sqrt{2} w_3 \end{pmatrix}$$

we clearly have

$$\hat{F}(0, 0, 0; 0) = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \quad \text{and} \quad J\hat{F}(0, 0, 0; 0) = B = \sqrt{2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & +1 \end{pmatrix}.$$

The eigenvector associated to $\lambda = 0$ is $v = (1, 0, 0)$ and since $B = B^T$ we have that the corresponding eigenvector to $\lambda = 0$ for this matrix is the same, namely $w^T = (1, 0, 0)$. Moreover, in view of the tangency conditions (3.9), a direct calculation shows

$$\hat{F}_\delta(0, 0, 0; 0) = (0, 0, 0); \quad D\hat{F}_\delta(0, 0, 0; 0) = \frac{1}{4} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Therefore $w^T \cdot \hat{F}_\delta(0, 0, 0; 0) = 0$ and $w^T \cdot D\hat{F}_\delta(0, 0, 0; 0)v = 1/4 \neq 0$. Finally we have to compute $w^T \cdot [D^2\hat{F}(0, 0, 0; 0)(v, v)]$ and we have to check that is different from zero. The computation of $[D^2\hat{F}(0, 0, 0; 0)(v, v)]$ gives the vector $(1/2, 0, 0)$ and thus $w^T \cdot [D^2\hat{F}(0, 0, 0; 0)(v, v)] \neq 0$. In view of *Sotomayor Theorem*, these three conditions imply that the original system (3.6) experiences a transcritical bifurcation at the equilibrium point $(0, 0, 0)$ as the parameter δ varies through the bifurcation value $\delta = 0$.

The presence of the parameter δ is only in the first component \hat{F}^1 of the vector field, for which, evaluating $\hat{F}_{w_1 w_1}^1(0, 0, 0; 0) = 1/2$ and $F_{\delta w_1}^1(0, 0, 0; 0) = 1/4$, we obtain the following the normal form (see for instance [4, Theorem 1.3]):

$$w'_1 = \frac{1}{4} \delta w_1 + \frac{1}{4} w_1^2 = \frac{1}{4} w_1 (w_1 + \delta).$$

Since $\delta < 0$, the trivial equilibrium $w_1 = 0$ is stable and $w_1 = -\delta > 0$ is unstable, and the same is true for the one-dimensional center manifold reduction given by

$$\begin{aligned} w'_1 = & \frac{1}{4} \{w_1 + \psi_2(w_1, \delta) - \psi_3(w_1, \delta)\}^2 \\ & + \frac{1}{4} \delta \{w_1 + \psi_2(w_1, \delta) - \psi_3(w_1, \delta)\}. \end{aligned} \tag{3.12} \quad \boxed{\text{eq:G1}}$$

As a consequence, there exists an heteroclinic connection between $w_1 = -\delta$ at $-\infty$ and $w_1 = 0$ at $+\infty$ for that equation. Then [9, Theorem 5.4, p. 159] implies that (3.6) is locally topologically equivalent to (3.12) augmented with the two linear equations

$$w'_2 = -\sqrt{2} w_2, \quad w'_3 = +\sqrt{2} w_3,$$

namely, system (3.10). The number of positive and negative eigenvalues is preserved, as well as the trajectories of the two dynamical systems. We have two unstable and one stable directions for the equilibrium point $(-\delta, 0, 0)$ and two stable directions and only one unstable at $(0, 0, 0)$. The two-dimensional unstable eigenspace of $(-\delta, 0, 0)$ and the two-dimensional stable eigenspace of $(0, 0, 0)$ intersect along the direction given by the vector $(1, 0, 0)$, namely the tangential direction of the center manifold in a neighborhood of the trivial equilibrium point. This implies the heteroclinic actually exists for (3.10). Back to the original variables, using (3.7) we have:

- 1) the point $(0, 0, 0)$ is mapped to $(u_+, 0, u_+)$,
- 2) the point $(-\delta, 0, 0)$ is mapped to $(u_-, 0, u_-)$,

and we have finally proved the existence of an heteroclinic orbit between these two points as stated in Theorem 3.1, thanks to the aforementioned local topological equivalence.

app:GSPT

APPENDIX A. GEOMETRIC SINGULAR PERTURBATION THEORY

A.1. Invariant Manifold Theorems. In this section we recall the main results about *Geometric Singular Perturbation Theory* developed in [5, 19]. Let us consider the vector field $X_\epsilon \times \{0\}$ defined in the following way:

$$\begin{cases} x' = \epsilon f(x, y, \epsilon) \\ y' = g(x, y, \epsilon) \\ \epsilon' = 0 \end{cases}$$

where $x \in \mathbb{R}^n$ are the slow variables, $y \in \mathbb{R}^k$ are the fast variables. The algebraic constraint $g(x, y, 0) = 0$ defines the slow manifold \mathcal{S} and correspondingly the two invertible branches \mathcal{S}^\pm . The following invariant manifold theorem describes the flow induced by $X_\epsilon \times \{0\}$ near $\mathcal{S}^\pm \times \{0\}$ for small ϵ .

existence

Theorem A.1. *Let M be a C^{r+1} manifold, $1 \leq r < \infty$. Let X_ϵ , $\epsilon \in (-\epsilon_0, \epsilon_0)$ be a C^r family of vector fields on M , and let \mathcal{S} be a C^r submanifold of M consisting entirely of equilibrium points of X_0 . Let k^s , k^c and k^u be fixed integers, and let $K \subset \mathcal{S}^\pm$ be a compact subset such that $QX_0(m)$ has k_s eigenvalues in the left half plane, k_c eigenvalues on the imaginary axis, and k_u in the right half plane, for all $m \in K$. Then:*

- 1) *There is a C^r center- stable manifold \mathcal{C}^s for $X_\epsilon \times 0$ near K . There is a C^r center- unstable manifold \mathcal{C}^u for $X_\epsilon \times 0$ near K . There is a C^r center manifold \mathcal{C} for $X_\epsilon \times 0$ near K .*
- 2) *There is a C^{r-1} family $\mathcal{F}^s = \{\Psi^s(p) : p \in \mathcal{C}^s\}$ of C^r stable manifolds for \mathcal{C}^s near K . If $p \in M \times \{\epsilon\}$, then $\Psi^s(p) \in M \times \{\epsilon\}$. Each manifold $\Psi^s(p)$ intersects \mathcal{C} transversally, in exactly one point. There is a*

C^{r-1} family $\mathcal{F}^u = \{\Psi^u(p) : p \in \mathcal{C}^u\}$ of C^r unstable manifolds for \mathcal{C}^u near K . If $p \in M \times \{\epsilon\}$, then $\Psi^u(p) \in M \times \{\epsilon\}$. Each manifold $\Psi^u(p)$ intersects \mathcal{C} transversally, in exactly one point.

- 3) Let $K_s < 0$ be larger then the real parts of the eigenvalues of $QX_0(m)$ in the left half plane, for all $m \in K$. Then, there is a constant C_s such that if $p \in \mathcal{C}^s$ and $q \in \Psi^s(p)$, then

$$d(p \cdot \bar{x}, q \cdot \bar{x}) \leq C_s e^{K_s \bar{x}} d(p, q)$$

for all $\bar{x} \geq 0$ such that $p \cdot [0, \bar{x}] \subset \mathcal{C}^s$. Let $K_u > 0$ be smaller than the real parts of the eigenvalues of $QX_0(m)$ in the right half plane, for all $m \in K$. Then there exists a constant C_u such that if $p \in \mathcal{C}^u$ and $q \in \Psi^u(p)$, then

$$d(p \cdot \bar{x}, q \cdot \bar{x}) \leq C_u e^{K_u \bar{x}} d(p, q)$$

for all $\bar{x} \leq 0$ such that $p \cdot [\bar{x}, 0] \subset \mathcal{C}^u$.

- 4) Let $S_H \subset S^\pm$ such that $QX_0(m)$ has not eigenvalue with zero real part. If $K \subset \mathcal{S}_H$, define for $(m, \epsilon) \in \mathcal{C}$,

$$\mathbf{F}_R(m) := \pi^S \left(\frac{\partial}{\partial \epsilon} \right) X_\epsilon(m)|_{\epsilon=0}$$

and

$$X_{\mathcal{C}}(m, \epsilon) := \begin{cases} \epsilon^{-1} X_\epsilon(m) \times \{0\}, & \text{if } \epsilon \neq 0 \\ \mathbf{F}_R(m) \times \{0\}, & \text{if } \epsilon = 0 \end{cases}$$

Then $X_{\mathcal{C}}$ is a C^{r-1} vector field on \mathcal{C} near $K \times \{0\}$.

A.2. Reduced System. The following lemma gives the reduced system in Theorem A.1 of the previous section in local coordinates in which \mathcal{S} appears as graph of a function (see [5]). From now on D_i , $i = 1, 2, 3$ denote the derivative with respect to slow variables D_x , fast variables D_y and D_ϵ respectively.

lemma3.1

Lemma A.2. Consider the system

$$\dot{x} = f(x, y, \epsilon), \quad \dot{y} = g(x, y, \epsilon),$$

defined for (x, y) in an open subset of $\mathbb{R}^n \times \mathbb{R}^k$, for ϵ near zero. Let $y = h(x)$ be a function such that

$$f(x, h(x), 0) \equiv 0, \quad \text{and} \quad g(x, h(x), 0) \equiv 0.$$

Suppose $(x_0, h(x_0)) \in \mathcal{S}_H$, so that the matrix

$$\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} \partial_x f(x_0, h(x_0), 0) & \partial_y f(x_0, h(x_0), 0) \\ \partial_x g(x_0, h(x_0), 0) & \partial_y g(x_0, h(x_0), 0) \end{pmatrix}$$

has rank k . Let $\nu = \partial_x h(x_0)$. Then the projection $\pi^S = \pi^S(x_0, u(x_0))$ is multiplication by the matrix

$$\mathbb{A} = \begin{pmatrix} I + \beta(\delta - \nu\beta)^{-1}\nu & -\beta(\delta - \nu\beta)^{-1} \\ \nu + \nu\beta(\delta - \nu\beta)^{-1}\nu & -\nu\beta(\delta - \nu\beta)^{-1} \end{pmatrix}$$

and the reduced system is given by:

$$\begin{pmatrix} \dot{x} \\ \dot{y} \end{pmatrix} = \mathbb{A} \begin{pmatrix} \partial_\epsilon f(x, h(x), 0) \\ \partial_\epsilon g(x, h(x), 0) \end{pmatrix}$$

A.3. Local theory near an equilibrium point of the reduced vector field. The next theorem explains how the critical points of the reduced vector field at $\epsilon = 0$ – denoted by \mathbf{F}_R – are related to the critical points of the original vector field X_ϵ in the regime of small $\epsilon > 0$.

Given $\epsilon \in (-\epsilon_0, \epsilon_0)$, let X_ϵ be a C^r family of vector fields on \mathbb{R}^3 and let \mathcal{S} be a C^r submanifold of \mathbb{R}^3 consisting entirely of equilibrium points of X_0 . Moreover, let $\mathbf{u} \in \mathcal{S}_H$ be an equilibrium point of the reduced vector field \mathbf{F}_R .

teo3.3 **Theorem A.3** (Theorem 12.1, [5]). *If $\lambda = 1$ is not an eigenvalue of $T\mathbf{F}_R(m)$, then there exists $\epsilon_1 > 0$ and a C^{r-1} family of points \mathbf{u}_ϵ with $\epsilon \in (-\epsilon_1, \epsilon_1)$ such that $\mathbf{u}^0 = \mathbf{u}$ and \mathbf{u}_ϵ is an equilibrium point of X_ϵ .*

Let M be a manifold and $\cdot t$ a flow on M . Given a subset $V \subset M$, we denote by

$$\begin{aligned} A^+(V) &:= \left\{ \mathbf{u} \in V : \overline{\mathbf{u} \cdot [0, +\infty)} \subset V \right\}, \\ A^-(V) &:= \left\{ \mathbf{u} \in V : \overline{\mathbf{u} \cdot (-\infty, 0]} \subset V \right\}, \\ I(V) &:= \left\{ \mathbf{u} \in V : \overline{\mathbf{u} \cdot (-\infty, \infty)} \subset V \right\}. \end{aligned}$$

The following result characterizes the local stable/unstable manifolds of normally hyperbolic invariant manifold of \mathbf{F}_R .

s-u-mxr **Theorem A.4** (Theorem 12.2, [5], Theorem 2.2, [19]). *Under the hypothesis of the previous Theorem, suppose $T\mathbf{F}_R(\mathbf{u}_0)$ has j^u eigenvalues in the right half plane, no eigenvalues on the imaginary axis and j^s eigenvalues in the left half plane. Suppose $Q\mathbf{F}_0(\mathbf{u}_0)$ has k^u eigenvalues in the right half plane and k^s eigenvalues in the left half plane.*

Then, there exists $\epsilon_1 > 0$ such that

- i. *there is a C^{r-1} -family of hyperbolic equilibrium points of \mathbf{F}_ϵ , denoted by $\{\mathbf{u}_\epsilon : \epsilon \in (-\epsilon_1, \epsilon_1)\}$, such that $\lim_{\epsilon \rightarrow 0} \mathbf{u}_\epsilon = \mathbf{u}_0$ and there is a family of neighborhoods of \mathbf{u}_0 , denoted by $\{U_\epsilon : \epsilon \in (-\epsilon_1, \epsilon_1)\}$, such that $I(U_\epsilon) = \{\mathbf{u}_\epsilon\}$ for any $\epsilon \neq 0$;*

- ii. *there is a C^{r-1} -families of $(j^u + k^u)$ -dimensional manifolds $\{W_\epsilon^u : \epsilon \in (-\epsilon_1, \epsilon_1)\}$ and $(j^s + k^s)$ -dimensional manifolds $\{\mathcal{W}_\epsilon^s : \epsilon \in (-\epsilon_1, \epsilon_1)\}$ such that*

$$A^-(U_\epsilon) = W_\epsilon^u \quad \text{and} \quad A^+(U_\epsilon) = \mathcal{W}_\epsilon^s \quad \forall \epsilon > 0;$$

- iii. *the local stable and unstable manifolds of \mathbf{u}_ϵ for $\epsilon > 0$ are given by*

$$N_\epsilon^s = \bigcup_{p \in \mathcal{W}_\epsilon^s} \Psi_\epsilon^s(p) \quad \text{and} \quad N_\epsilon^u = \bigcup_{p \in \mathcal{W}_\epsilon^u} \Psi_\epsilon^u(p),$$

where $\{\Psi_\epsilon^s(\mathbf{u}) : \epsilon \in (-\epsilon_1, \epsilon_1), \mathbf{u} \in \mathcal{W}_\epsilon^s\}$ is a C^{r-1} -family of k_1 -dimensional manifolds such that $\{\Psi_\epsilon^s(m) : m \in \mathcal{W}_\epsilon^s\}$ for each $\epsilon > 0$ is a positively invariant family of manifolds (the same for $\Psi_\epsilon^u(m)$).

theo:a5

Theorem A.5 (Theorem 3.1, [19]). *Let the manifolds N_1 and N_2 satisfy the assumptions of Theorem A.4. Assume that the manifolds*

$$N_1^u := \bigcup_{p \in \mathcal{W}_1^u} \Psi^u(p), \quad N_2^s := \bigcup_{p \in \mathcal{W}_2^s} \Psi^s(p)$$

intersect transversally along the singular heteroclinic orbit. Then, there exists $\epsilon_1 > 0$ such that there exists a transversal heteroclinic orbit of the singularly perturbed system (1.1) in [19] connecting the manifolds $N_{1,\epsilon}$ and $N_{2,\epsilon}$ for $0 < \epsilon < \epsilon_1$.

Finally, the last needed result concerns sufficient (transversality) conditions needed to obtain the connections between the two branches of the involved invariant manifold.

teo3.12

Theorem A.6 (Theorem 4.1, [19]). *Let the manifolds N_1 and N_2 satisfy the assumption of Theorem A.4. Let $\phi(W_1^u)$ and $\phi(W_2^s)$ denote the x -coordinates of the manifolds \mathcal{W}_1^u and \mathcal{W}_2^s . Then the manifolds N_1^u and N_2^s intersect transversally in the points of the heteroclinic orbit $(x_0, y_0(x))$, if and only if there exist exactly $d - 1$ linearly independent solutions $\xi \in T_{x_0}\phi(\mathcal{W}_1^u) \cap T_{x_0}\phi(\mathcal{W}_2^s)$ of the equation*

$$(M, \xi) = 0, \tag{A.1} \quad \text{eq:M}$$

where $M \in \mathbb{R}^n$ is defined by

$$M := \int_{\mathbb{R}} \psi(\xi) \cdot \partial_x g(x_0, y_0(\xi)) d\xi.$$

The function ψ is the unique (up to a scalar multiple) bounded solution of the adjoint equation

$$\psi' = -\partial_y g(x_0, y_0(x))^T \psi.$$

REFERENCES

- [1] Buet C., Després B.; Asymptotic analysis of fluid models for the coupling of radiation and hydrodynamics, *J. Quant. Spectrosc. Radiat. Transf.* **85**:3-4 (2004) 385–418.
- [2] Corli A., Rohde C.; Singular limits for a parabolic-elliptic regularization of scalar conservation laws, *J. Differ. Equations* **253**:5 (2012) 1399–1421.
- [3] Coulombel J.-F., Goudon T., Lafitte P., Lin C.; Analysis of large amplitude shock profiles for non-equilibrium radiative hydrodynamics: formation of Zeldovich spikes, *Shock Waves* **22**:3 (2012) 181–197.
- [4] Faye G.; “An introduction to bifurcation theory”, online, October 6th, 2011
- [5] Fenichel N.; Geometric singular perturbation theory for ordinary differential equations, *J. Differ. Equations* **31**:1 (1979) 53–98.
- [6] Godillon-Lafitte P., Goudon T.; A coupled model for radiative transfer: Doppler effects, equilibrium and non equilibrium diffusion asymptotics, *Multiscale Model. Sim.* **4**:4 (2005) 1245–1279.
- [7] Hamer K.; Nonlinear effects on the propagation of sound waves in a radiating gas, *Quart. J. Mech. Appl. Math.* **24** (1971) 155–168.
- [8] Kawashima S., Nishibata S.; Shock waves for a model of the radiating gas, *SIAM J. Math. Anal.* **30**:1 (1999) 95–117.
- [9] Kuznetsov Y.A.; “Elements of applied bifurcation theory”. Third edition. Applied Mathematical Sciences, 112, *Springer-Verlag, New York*, 2004.
- [10] Lattanzio C., Marcati P.; Global well-posedness and relaxation limits of a model for radiating gas, *J. Differential Equations* **190**:2 (2003) 439–465.
- [11] Lattanzio C., Mascia C., Serre D.; Shock waves for radiative hyperbolic-elliptic systems, *Indiana Univ. Math. J.* **56**:5 (2007) 2601–2640.
- [12] Lattanzio C., Mascia C., Serre D.; Nonlinear hyperbolic-elliptic coupled systems arising in radiation dynamics, in: S. Benzoni-Gavage, D. Serre (Eds.), *Hyperbolic Problems: Theory, Numerics, Applications (Lyon, July 17-21, 2006)*, Springer-Verlag, Boston, Berlin, Heidelberg, pp. 661–669, 2008.
- [13] Lin C., Coulombel J.-F., Goudon T.; Shock profiles for non-equilibrium radiating gases, *Physica D* **218**:1 (2006) 83–94.
- [14] Lowrie R.B., Morel J.E., Hittinger J.A.; The coupling of radiation and hydrodynamics, *Astrophys. J.* **521**:1 (1999) 432–450.
- [15] Mascia C.; Small, medium and large shock waves for radiative Euler equations, *Physica D* **245**:1 (2013) 46–56.
- [16] Mihalas D., Mihalas B.W., “Foundations of Radiation Hydrodynamics”, *Oxford University Press, New York*, 1984. Reprinted by Dover Publications, New York, 1999.
- [17] Perko L.; “Differential equations and dynamical systems”. Third edition. Texts in Applied Mathematics, 7. *Springer-Verlag, New York*, 2001.
- [18] Schochet S., Tadmor E.; The regularized Chapman–Enskog expansion for scalar conservation laws, *Arch. Ration. Mech. Anal.* **119**:2 (1992), 95–107.
- [19] Szmolyan P.; Transversal heteroclinic and homoclinic orbits in singular perturbation problem, *J. Differ. Equations* **92**:2 (1991) 252–281.
- [20] Xuan M., Hui T., Jin J.; Global asymptotic toward the rarefaction waves for a parabolic-elliptic system related to the Camassa–Holm shallow water equation, *Acta Math. Sci.* **29B**:2 (2009) 371–390.
- [21] Zel’dovich Y.B., Raizer Y.P.; “Physics of Shock Waves and High-Temperature Hydrodynamic Phenomena”, Academic Press, New York 1967. Reprinted by Dover Publications, New York, 2002.

(Giada Cianfarani Carnevale) DIPARTIMENTO DI INGEGNERIA E SCIENZE DELL'INFORMAZIONE
E MATEMATICA, UNIVERSITÀ DEGLI STUDI DELL'AQUILA (ITALY)

E-mail address: `giada.cianfaranicarnevale@graduate.univaq.it`

(Corrado Lattanzio) DIPARTIMENTO DI INGEGNERIA E SCIENZE DELL'INFORMAZIONE E
MATEMATICA, UNIVERSITÀ DEGLI STUDI DELL'AQUILA (ITALY)

E-mail address: `corrado@univaq.it`

(Corrado Mascia) DIPARTIMENTO DI MATEMATICA "GUIDO CASTELNUOVO", SAPIENZA
UNIVERSITÀ DI ROMA (ITALY)

E-mail address: `corrado.mascia@uniroma1.it`