

STATIONARY NON EQUILIBRIUM STATES FROM A MICROSCOPIC AND A MACROSCOPIC POINT OF VIEW

Davide Gabrielli

University of L'Aquila

1 July 2014

GGI Firenze

References


- **L. Bertini; A. De Sole; D. G. ; G. Jona-Lasinio; C. Landim**
Stochastic interacting particle systems out of equilibrium J.
Stat. Mech. (2007)
- **D. G.** From combinatorics to large deviations for the
invariant measures of some multiclass particle systems
Markov Processes Relat. (2008)
- **L. Bertini; D. G.; G. Jona-Lasinio; C. Landim**
Thermodynamic transformations of nonequilibrium states
J. Stat. Phys. (2012)

SNS: Microscopic description

Lattice: Λ_N

Configuration of particles: $\eta \in \{0, 1\}^{\Lambda_N}$ or $\eta \in \mathbb{N}^{\Lambda_N}$

$\eta_t(x)$ = number of particles at $x \in \Lambda_N$ at time t



SNS: Microscopic description

- Stochastic Markovian dynamics
- $r(\eta, \eta')$ = rate of jump from configuration η to configuration η'
- η' = local perturbation of η
- $\mu_N(\eta)$ = invariant measure of the process, probability measure on the state space

$$\mu_N(\eta) \sum_{\eta'} r(\eta, \eta') = \sum_{\eta'} \mu_N(\eta') r(\eta', \eta)$$

$\mu_N \implies$ MICROSCOPIC description of the SNS

SNS: Macroscopic description


- $\eta \implies \pi_N(\eta)$ Empirical measure (positive measure on $[0, 1]$)

$$\pi_N(\eta) = \frac{1}{N} \sum_{x \in \Lambda_N} \eta(x) \delta_x$$

$\delta_x =$ delta measure (Dirac) at $x \in [0, 1]$; since $x \in \Lambda_N$ we have $x = \frac{i}{N}$, $i \in \mathbb{N}$. Given $f : [0, 1] \rightarrow \mathbb{R}$

$$\int_{[0,1]} f d\pi_N = \frac{1}{N} \sum_{x \in \Lambda_N} \eta(x) f(x)$$

X = delta measure



SNS: Macroscopic description

When η is distributed according to μ_N and N is large

LAW OF LARGE NUMBERS

$$\pi_N \rightarrow \bar{\rho}(x)dx$$

This means

$$\int_{[0,1]} f d\pi_N \rightarrow \int_{[0,1]} f(x) \bar{\rho}(x) dx$$

$\bar{\rho}(x)$ = typical density profile of the SNS

SNS: Macroscopic description

When η is distributed according to μ_N and N is large, a refinement of the law of large numbers

LARGE DEVIATIONS

$$\mathbb{P}\left(\pi_N(\eta) \sim \rho(x)dx\right) \simeq e^{-NV(\rho)}$$

V = Large deviations rate function


$V \implies$ MACROSCOPIC DESCRIPTION OF THE SNS

V contains less information than μ_N but is easier to compute and is independent from microscopic details of the dynamics

Example: Equilibrium SEP

- Equilibrium: $C_L = C_R = C$; $A_L = A_R = A$
- Microscopic state: product of Bernoulli measures of parameter $p = \frac{C}{A+C}$

$$\mu_N(\eta) = \prod_{x \in \Lambda_N} p^{\eta(x)} (1-p)^{1-\eta(x)}$$



Example: Equilibrium SEP

MACROSCOPIC DESCRIPTION

$$\begin{aligned} \mathbb{P}\left(\pi_N(\eta) \sim \rho(x)dx\right) &= \sum_{\{\eta, : \pi_N(\eta) \sim \rho(x)dx\}} \mu_N(\eta) \\ &= \sum_{\{\eta, : \pi_N(\eta) \sim \rho(x)dx\}} e^{-N \left(\int_{[0,1]} d\pi_N(\eta) \log \frac{1-p}{p} - \log(1-p) \right)} \end{aligned}$$

Using the combinatorial estimate

$$\left| \{\eta, : \pi_N(\eta) \sim \rho(x)dx\} \right| \simeq e^{-N \int_0^1 \rho(x) \log \rho(x) + (1-\rho(x)) \log(1-\rho(x)) dx}$$

$$V(\rho) = \int_0^1 \rho(x) \log \frac{\rho(x)}{p} + (1 - \rho(x)) \log \frac{(1-\rho(x))}{(1-p)} dx$$

Contraction

Average number of particles

$$\frac{1}{N} \sum_i \eta(i) = \int_{[0,1]} d\pi_N(\eta)$$

satisfies LDP

$$\mathbb{P} \left(\frac{1}{N} \sum_i \eta(i) \sim y \right) \simeq e^{-NJ(y)}$$

BY CONTRACTION

$$J(y) = \inf_{\{\rho: \int_0^1 \rho(x) dx = y\}} V(\rho)$$

Relative entropy

Relative entropy of the probability measure μ_N^2 with respect to μ_N^1

$$H\left(\mu_N^2 \middle| \mu_N^1\right) = \sum_{\eta} \mu_N^2(\eta) \log \frac{\mu_N^2(\eta)}{\mu_N^1(\eta)}$$

$H \geq 0$, not symmetric!!

Density of relative entropy

$$h = \lim_{N \rightarrow +\infty} \frac{1}{N} H\left(\mu_N^2 \middle| \mu_N^1\right)$$

An example

$\mu_N^1(\eta) = \prod_{x \in \Lambda_N} p^{\eta(x)} (1-p)^{1-\eta(x)}$, product of Bernoulli measures of parameter p

$\mu_N^2(\eta) = \prod_{x \in \Lambda_N} \rho(x)^{\eta(x)} (1-\rho(x))^{1-\eta(x)}$, slowly varying product of Bernoulli measures associated to the density profile $\rho(x)$

$$\begin{aligned} & \frac{1}{N} H(\mu_N^2 | \mu_N^1) \\ &= \sum_{\eta} \mu_N^2(\eta) \left(\frac{1}{N} \sum_{x \in \Lambda_N} \eta(x) \log \frac{\rho(x)}{p} + (1 - \eta(x)) \log \frac{(1 - \rho(x))}{(1 - p)} \right) \\ &= \frac{1}{N} \sum_{x \in \Lambda_N} \rho(x) \log \frac{\rho(x)}{p} + (1 - \rho(x)) \log \frac{(1 - \rho(x))}{(1 - p)} \end{aligned}$$

Riemann sums, convergence when $N \rightarrow +\infty$ to $V(\rho)$

From microscopic to MACROSCOPIC


- Driving parameters (λ, E)
- $\lambda \implies$ rates of injection and annihilation at the boundary
- $E \implies$ external field driving the particles on the bulk
- $\mu_N^{\lambda, E} \implies$ corresponding invariant measure
- $\bar{\rho}_{\lambda, E} \implies$ corresponding typical density profile
- $V_{\lambda, E}(\rho) \implies$ corresponding LD rate function

$$V_{\lambda_1, E_1}(\bar{\rho}_{\lambda_2, E_2}) = \lim_{N \rightarrow +\infty} \frac{1}{N} H\left(\mu_N^{\lambda_2, E_2} \middle| \mu_N^{\lambda_1, E_1}\right)$$

From microscopic to MACROSCOPIC


- This relation between relative entropy and LD rate function can be easily verified for the boundary driven Zero Range Process
- It is true also for boundary driven SEP; proof based on matrix representation of μ_N
- In general the computation of V through relative entropy is difficult
- An alternative powerful approach to compute V is the dynamic variational one of the Macroscopic Fluctuation Theory

Boundary driven TASEP: a microscopic view



Boundary driven TASEP: a microscopic view

- **Duchi E., Schaeffer G** A combinatorial approach to jumping particles, J. Comb. Theory A (2005)



Boundary driven TASEP: a microscopic view

- $\eta \implies$ configuration of particles above
- $\xi \implies$ configuration of particles below
- $(\eta, \xi) \implies$ full configuration of particles
- Stochastic Markov dynamics for (η, ξ)
- Observing just $\eta \implies$ still Markov and boundary driven TASEP
- $\nu_N(\eta, \xi) \implies$ invariant measure for the joint dynamics, it has a combinatorial representation

$$\mu_N(\eta) = \sum_{\xi} \nu_N(\eta, \xi)$$

Boundary driven TASEP: a microscopic view

Complete configurations

$$E(x) = \sum_{y \leq x} (\eta(y) + \xi(y)) - Nx - 1$$

(η, ξ) is a **complete configuration** if

$$\begin{cases} E(x) \geq 0 \\ E(1) = 0 \end{cases}$$

ν_N is concentrated on complete configurations

(η, ξ) complete $\implies N_1(\eta, \xi), N_2(\eta, \xi)$

$$\nu_N(\eta, \xi) = \frac{1}{Z_N} A^{N_1(\eta, \xi)} C^{N_2(\eta, \xi)}$$

Special case $A = C = 1 \implies \nu_N$ uniform measure on complete configurations

Boundary driven TASEP: a macroscopic view

Joint Large deviations

$$\mathbb{P}\left(\left(\pi_N(\eta), \pi_N(\xi)\right) \sim \left(\rho(x), f(x)\right)\right) \simeq e^{-N\mathcal{G}(\rho, f)}$$

Contraction principle

$$\mathbb{P}\left(\pi_N(\eta) \sim \rho(x)\right) \simeq e^{-NV(\rho)}$$

$$V(\rho) = \inf_f \mathcal{G}(\rho, f)$$

Boundary driven TASEP: a macroscopic view

Complete density profiles

$$\mathcal{E}(x) = \int_0^x (\rho(y) + f(y)) dy - x$$

The pair (ρ, f) is a **complete density profile** if

$$\begin{cases} \mathcal{E}(x) \geq 0 \\ \mathcal{E}(1) = 0 \end{cases}$$

When $C = A = 1$ since ν_N is uniform on complete configurations a classic simple computation gives

$$\mathcal{G}(\rho, f) = \int_0^1 \left[h_{\frac{1}{2}}(\rho(x)) + h_{\frac{1}{2}}(f(x)) \right] dx$$

if (ρ, f) is complete; here

$$h_p(\alpha) = \alpha \log \frac{\alpha}{p} + (1 - \alpha) \log \frac{(1 - \alpha)}{1 - p}$$

Boundary driven TASEP: a macroscopic view

$$V(\rho) = \inf_{f: (\rho, f) \in \mathcal{C}} \int_0^1 \left[h_{\frac{1}{2}}(\rho(x)) + h_{\frac{1}{2}}(f(x)) \right] dx$$

To be compared with **B. Derrida, J.L. Lebowitz, E.R. Speer**
Exact large deviation functional of a stationary open driven
diffusive system: the asymmetric exclusion process *J. Stat.*
Phys. (2003)

$$V(\rho) = \sup_f \int_0^1 \left\{ \rho(x) \log [\rho(x)(1 - f(x))] \right. \\ \left. + (1 - \rho(x)) \log [(1 - \rho(x))f(x)] \right\} dx + \log 4$$

where $f(0) = 1$, $f(1) = 0$ and f is monotone


Boundary driven TASEP: a macroscopic view

Both variational problems have the same minimizer


$$f_\rho(x) = CE\left(\int_0^x (1 - \rho(y)) dy\right)$$

$$V(\rho) = \mathcal{G}(\rho, f_\rho)$$


See [Bahadoran C.](#) A quasi-potential for conservation laws with boundary conditions arXiv:1010.3624 for a dynamic variational approach, using MFT



2-class TASEP



The invariant measure



Collapsing particles

$$(\tilde{\eta}_1, \tilde{\eta}_T) : \sum_x \tilde{\eta}_1(x) \leq \sum_x \tilde{\eta}_T(x) \implies (\eta_1, \eta_T) = \mathcal{C}[(\tilde{\eta}_1, \tilde{\eta}_T)]$$

Flux across bond $(x, x + 1)$

$$J(x) = \sup_y \left[\sum_{z \in [y, x]} \tilde{\eta}_1(z) - \tilde{\eta}_T(z) \right]_+$$

Collapsing measures

$$(\tilde{\rho}_1, \tilde{\rho}_T) : \int_{\mathbb{S}^1} d\tilde{\rho}_1 \leq \int_{\mathbb{S}^1} d\tilde{\rho}_T \implies (\rho_1, \rho_T) = \mathcal{C}[(\tilde{\rho}_1, \tilde{\rho}_T)]$$

Definition

$$\int_{(a,b]} d\rho_1 = \int_{(a,b]} d\tilde{\rho}_1 + J(a) - J(b)$$

where

$$J(x) := \sup_y \left[\int_{(y,x]} d\tilde{\rho}_1 - \int_{(y,x]} d\tilde{\rho}_2 \right]_+$$

Collapsing measures



Large deviations

LD for the $(\tilde{\eta}_1, \tilde{\eta}_T)$ variables

$$\tilde{V}(\tilde{\rho}_1, \tilde{\rho}_T) = \int_{\mathbb{S}^1} [h_{m_1}(\tilde{\rho}_1) + h_{m_2}(\tilde{\rho}_T)] dx$$

LD for the SNS (not convex!)

$$\begin{aligned} V(\rho_1, \rho_T) &= \inf_{\{(\tilde{\rho}_1, \tilde{\rho}_T) : \mathcal{C}[(\tilde{\rho}_1, \tilde{\rho}_T)] = (\rho_1, \rho_T)\}} \tilde{V}(\tilde{\rho}_1, \tilde{\rho}_T) \\ &= \int_{\mathbb{S}^1} [h_{m_1}(\hat{\rho}_1) + h_{m_2}(\rho_T)] dx \end{aligned}$$

On any (a, b) where $\rho_1 = \rho_T$

$$\int_a^x \hat{\rho}_1(y) dy = CE \left[\int_a^x \rho_1(y) dy \right]$$